

A note on partial isometries on pseudo-Hilbert spaces

Păstorel Gaşpar

Loredana Ciurdariu

Abstract

The aim of this paper is to show that two accessible subspaces in the Loynes $\mathcal{Z}$ - space $\mathcal{H}$ are the initial and final space of a partial gramian isometry, respectively if the norm of the difference of the associated gramian selfadjoint projections is strictly less than 1.¹

1 Introduction

Generalizing the concept of pre-Hilbert or Hilbert space, R.M. Loynes introduced in [5] the VE - spaces or VH - spaces respectively. A VH - space is characterized in [6] by the fact that the inner product takes values in a suitable ordered topological vector (admissible) space $\mathcal{Z}$, thus being also called *Loynes $\mathcal{Z}$ - spaces*. Many authors used these spaces in the study of abstract stochastic processes (see [7], [1], [11], [12]). In [11] these spaces are referred to as *pseudo-Hilbert spaces*. Spectral theory for some classes of operators on such spaces was developed initially by Loynes himself ([5], [6]) and later by the authors in [3], respectively [2] and by A. Gheondea and B. E. Ugurcan in [4].

In what follows $\mathcal{H}, \mathcal{K}$ will denote two pseudo-Hilbert spaces over the same admissible space $\mathcal{Z}$ and $\mathcal{L}(\mathcal{H}, \mathcal{K})$ the space of all linear operators from $\mathcal{H}$ to $\mathcal{K}$.

Recall that an operator $T \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ is *bounded*, if there exists a constant $M > 0$ such that

$$[Th, Th]_{\mathcal{K}} \leq M^2[h, h]_{\mathcal{H}}, \quad h \in \mathcal{H}, \quad (1)$$

where $[\cdot, \cdot]_{\mathcal{K}}$ is the inner product (also referred to as *gramian*) of the Loynes $\mathcal{Z}$ - space $\mathcal{K}$, while “ $\leq$ ” means the order in $\mathcal{Z}$. We shall denote that by

¹AMS Subject Classification: 46C50, 47B37

Keywords: pseudo-Hilbert spaces, partial gramian isometry, gramian selfadjoint projection

$T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$. As usually, for $\mathcal{H} = \mathcal{K}$, we use the notations $\mathcal{L}(\mathcal{H})$ and $\mathcal{B}(\mathcal{H})$ respectively. Moreover $\mathcal{B}(\mathcal{H}, \mathcal{K})$ is a Banach space (algebra if $\mathcal{H} = \mathcal{K}$) with the norm defined by

$$\|T\| = \|T\|_{\mathcal{B}(\mathcal{H}, \mathcal{K})} = \inf\{M : (1) \text{ holds } \}.$$

The operators $T \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ for which $\|T\| \leq 1$ will be called *gramian contractions*.

The adjoint T^* of an operator $T \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ and the gramian orthogonal complement $\mathcal{M}^\perp$ of a subspace $\mathcal{M}$ of $\mathcal{H}$ will be defined (if they exist) analogously as in the Hilbert space case, but with respect to the inner products of $\mathcal{H}$ and $\mathcal{K}$.

By $\mathcal{L}^*(\mathcal{H}, \mathcal{K})$, $\mathcal{B}^*(\mathcal{H}, \mathcal{K})$ will be denoted the set of all adjointable elements of $\mathcal{L}(\mathcal{H}, \mathcal{K})$ and $\mathcal{B}(\mathcal{H}, \mathcal{K})$, respectively, whereas $P_{\mathcal{M}}$ denotes the gramian selfadjoint projection associated to the complementable (accessible) subspace $\mathcal{M}$ of $\mathcal{H}$.

We also remark that $\mathcal{L}^*(\mathcal{H}, \mathcal{K}) \cap \mathcal{B}(\mathcal{H}, \mathcal{K}) = \mathcal{B}^*(\mathcal{H}, \mathcal{K})$ and $\mathcal{B}^*(\mathcal{H})$ is a C^* -algebra.

$T \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ is called a *gramian isometry* (*gramian co-isometry*) if $T \in \mathcal{B}^*(\mathcal{H}, \mathcal{K})$ and $T^*T = I_{\mathcal{H}}$ ($TT^* = I_{\mathcal{K}}$, respectively) and T is *gramian unitary* if it is simultaneously a gramian isometry and a gramian co-isometry.

If a gramian contraction T is adjointable, then T^* is a gramian contraction too (see [5], [6]). Familiar examples of adjointable contractions are self-adjoint gramian projections and gramian partial isometries, the latter containing two remarkable subclasses: that of gramian isometries and of gramian co-isometries. The latter classes were studied by the first author in [3], where also a geometric proof of the existence of the gramian co-isometric extension of a gramian adjointable contraction is given.

In what follows some definitions and results from [3] are needed.

Definition 1.1. *Let $\mathcal{H}$ and $\mathcal{K}$ be two Loynes $\mathbb{Z}$ -spaces. A linear operator $T \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ is a *partial gramian isometry*, if its kernel $\mathcal{N}(T)$ and its range $\mathcal{R}(T)$ are accessible (i.e. they have gramian orthogonal complements) in $\mathcal{H}$ and $\mathcal{K}$, respectively and from $\mathcal{N}(T)^\perp$ to $\mathcal{R}(T)$ it preserves the gramian (is gramian unitary). The spaces $\mathcal{M} := \mathcal{N}(T)^\perp$ and $\mathcal{R}(T)$ are called the *initial* and the *final space* of T , respectively. The set of all partial gramian isometries from $\mathcal{H}$ to $\mathcal{K}$ will be denoted by $\mathcal{PJ}(\mathcal{H}, \mathcal{K})$.*

It can be easily seen, that $\mathcal{PJ}(\mathcal{H}, \mathcal{K}) \subset \mathcal{B}^*(\mathcal{H}, \mathcal{K})$. Observe that if $\mathcal{N}(T) = 0$, then T is simply a gramian isometry.

Proposition 1.1. *If $T \in \mathcal{PJ}(\mathcal{H}, \mathcal{K})$, then T^*T and TT^* are gramian self-adjoint projections for which the following hold:*

- (i) $T^*T = P_{\mathcal{M}(T)}$;
- (ii) $P_{\mathcal{R}(T)} = TT^*$;
- (iii) If $\mathcal{H} = \mathcal{K}$ then T is a partial isometry in $\mathcal{B}^*(\mathcal{H})$ as a C^* -algebra.

Proposition 1.2. *For $T \in \mathcal{L}(\mathcal{H}, \mathcal{K})$, the following are equivalent:*

- (i) $T \in \mathcal{PJ}(\mathcal{H}, \mathcal{K})$;
- (ii) $T \in \mathcal{B}^*(\mathcal{H}, \mathcal{K})$ and T^*T is a gramian self-adjoint projection on $\mathcal{H}$;
- (iii) $T \in \mathcal{B}^*(\mathcal{H}, \mathcal{K})$ and TT^* is a gramian self-adjoint projection on $\mathcal{K}$;
- (iv) $T \in \mathcal{L}^*(\mathcal{H}, \mathcal{K})$ and $T^* \in \mathcal{PJ}(\mathcal{K}, \mathcal{H})$.

It is obvious that any gramian isometry or gramian co-isometry is a partial gramian isometry.

2 The result

Focusing on the case $\mathcal{H} = \mathcal{K}$ and taking $T \in \mathcal{PJ}(\mathcal{H})$, then the operators T^*T and TT^* will be two gramian self-adjoint projections in $\mathcal{B}^*(\mathcal{H})$. It is thus interesting, as in the case of Hilbert space (see [10, pp. 266,267]), to find a sufficient condition on two gramian self-adjoint projections P and Q in order to have their ranges as initial and final space of a certain partial gramian isometry. Indeed the following assertion holds.

Theorem 2.1. *If P and Q are gramian self-adjoint projections and*

$$\|P - Q\| < 1, \tag{2}$$

then there exists $T \in \mathcal{PJ}(\mathcal{H})$ such that $P = T^*T$ and $Q = TT^*$.

Proof. Denote $A = I + P(Q - P)P$. Since $\|I - A\| = \|P(Q - P)P\| \leq \|P - Q\| < 1$, by using that $\mathcal{B}^*(\mathcal{H})$ is a Banach algebra, it results that A is invertible with a bounded inverse. On the other hand the operator A is positive. Indeed

$$\begin{aligned} [Ah, h] &= [h, h] + [P(Q - P)Ph, h] = [h, h] + [QPh, h] - [P^3h, h] \\ &= [(I - P)h, h] + [QPh, Ph] \geq 0, \end{aligned}$$

where we used the fact that $I - P$ and Q are gramian self-adjoint projections. In this situation, there exists the square root of A , which is also invertible.

The operator $T := QA^{-1/2}P$ satisfies the requirements of the statement. Indeed we have $T^* = PA^{-1/2}Q$ and further on $PT^* = T^* = A^{-1/2}PQ$. Since $PA = AP$, we infer that $PA^{1/2} = A^{1/2}P$ which implies $A^{-1/2}P = PA^{-1/2}$. Further we get $TP = T = QPA^{-1/2}$. Taking into account that $PA = PQP$ we infer

$$T^*T = A^{-1/2}PQQPA^{-1/2} = A^{-1/2}PQPA^{-1/2} = A^{-1/2}PAA^{-1/2} = P.$$

Using Proposition 1.2 we infer that T is a partial gramian isometry and $TT^* = P_{\mathcal{R}(T)}$. But, the calculation of TT^* leads us to the equalities

$$TT^* = QA^{-1/2}PPA^{-1/2}Q = QA^{-1}PQ,$$

which imply $\mathcal{R}(T) = \mathcal{R}(TT^*) \subset \mathcal{R}(Q)$, i.e. $TT^* \leq Q$. Now, let us show that $I - TT^* \leq I - Q$. Let $h \in (I - TT^*)\mathcal{H}$. Then the next implications hold

$$\begin{aligned} h \in (I - TT^*)\mathcal{H} &\Rightarrow h = (I - TT^*)h \Rightarrow TT^*h = 0 \\ &\Rightarrow R^*h \in \mathcal{N}(T) \cap \mathcal{R}(T^*) = \{0\} \Rightarrow T^*h = 0 \\ &\Rightarrow PA^{-1/2}Qh = 0 \Rightarrow PQh = 0 \Rightarrow (Q - P)Qh = Qh \\ &\Rightarrow Qh = 0 \Rightarrow (I - Q)h = h \\ &\Rightarrow h \in \mathcal{R}(I - Q). \end{aligned}$$

This shows that $\mathcal{R}(I - TT^*) \subset \mathcal{R}(I - Q)$, which indicates that $I - TT^* \leq I - Q$. Hence the equality $TT^* = Q$ holds. $\square$

Remark 2.1. Our theorem can be applied in perturbation theory to treat the variation of the spectral measure of gramian selfadjoint operators on pseudo-Hilbert spaces in a limit taking process. For the Hilbert space case see [10] no. 135.

Remark 2.2. Our theorem states that (2) is a sufficient condition on the two gramian selfadjoint projections P and Q in order to determine the initial and final space of a partial isometry. This condition isn't however necessary, as the following example shows. For V a gramian (non-unitary) isometry on $\mathcal{H}$ we have that $V^*V - VV^*$, being a gramian selfadjoint projection, has norm equal to 1.

It would therefore be interesting to find a weaker condition that would still be sufficient.

Remark 2.3. Our definition of the partial isometry on the pseudo-Hilbert space $\mathcal{H}$ as well as the statement of our result being given in the C^* -algebra $\mathcal{B}^*(\mathcal{H})$ let us observe that following [8] or [9] it is possible to define and characterize the notion of a partial isometry in the Banach algebra $\mathcal{B}(\mathcal{H})$ ($\square$)

$\mathcal{B}^*(\mathcal{H})$). It is then naturally to ask if there exist such partial isometries in $\mathcal{B}(\mathcal{H})$ which are not in $\mathcal{B}^*(\mathcal{H})$ and if this would be the case, would an analogue of Theorem 2.1 hold in $\mathcal{B}(\mathcal{H})$?

References

- [1] S.A. Chobanyan, A. Weron, *Banach-space-valued stationary processes and their linear prediction*, Dissertationes Math., 125, (1975), 1-45.
- [2] L. Ciurdariu, A. Crăciunescu, *On Spectral Representation of Gramian Normal Operators on Pseudo-Hilbert Spaces*, Anal. Univ. de Vest Timișoara, Vol. XLV, (1), 2007, 131-149.
- [3] P. Gașpar, *Partial isometries on Loynes spaces*, Anal. Univ. de Vest Timișoara, Vol. XL, (2), 2002, 31-47.
- [4] A. Gheondea, B. E. Ugurcan, *On two equivalent dilation theorems in VH - spaces*, Complex Analysis and Operator Theory, vol. 6, (3), 2012, 625-650.
- [5] R.M. Loynes, *Linear operators in VH -spaces*, Trans. American Math. Soc., 116, (1965), 167-180.
- [6] R.M. Loynes, *On generalized positive definite functions*, Proc. London Math. Soc., 3, (1965), 373-384.
- [7] R.M. Loynes, *On a generalization of second-order stationarity*, Proc. London Math. Soc., 3, (1965), 385-398.
- [8] M. Mbekhta, *Partial isometries and generalized inverses*, Acta. Sci. Math. (Szeged) **70** (2004), 767-781.
- [9] D. Mosič, D. S. Djordjevič, *Partial isometries and EP elements in Banach algebras*, preprint, http://operator.pmf.ni.ac.rs/licne_prezentacije/DDjordjevic/publications/Partial-EP-Banach.pdf
- [10] F. Riesz, B.Sz.-Nagy, *Leçons d'analyse fonctionnelle*, 4-ème Edition, Gauthier-Villars Paris and Akadémiai Kiadó Budapest, 1965.
- [11] A. Weron, S.A. Chobanyan, *Stochastic Processes on pseudo-Hilbert spaces(russian)*, Bull. Acad. Pol., Ser. Math. Phys., tom XX1, 9, (1973), 847-854.

- [12] A. Weron, *Prediction theory in Banach spaces*, Proc. of Winter School on Probability, Karpacz, Springer Verlag, London, 1975, 207-228.

Păstorel Gaşpar, “Aurel Vlaicu” University, Arad, e-mail: pastogaspar@yahoo.com
Loredana Ciurdariu, “Politehnica” University, Timișoara